

# WATCAT: A Tale of Wide-Angle Tailed Radio Galaxies

V. Missaglia<sup>1</sup>, F. Massaro<sup>2,3,4,5</sup>, A. Capetti<sup>4</sup>, M. Paolillo<sup>6,7</sup>, R.P. Kraft<sup>1</sup>, R.D. Baldi<sup>8</sup>, and A. Paggi<sup>2,3,4</sup>

<sup>1</sup> Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA

<sup>2</sup> Dipartimento di Fisica, Università degli Studi di Torino, via Pietro Giuria 1, 10125 Torino, Italy

<sup>3</sup> Istituto Nazionale di Fisica Nucleare, Sezione di Torino, I-10125 Torino, Italy

<sup>4</sup> INAF - Osservatorio Astrofisico di Torino, via Osservatorio 20, 10025 Pino Torinese, Italy

<sup>5</sup> Consorzio Interuniversitario per la Fisica Spaziale (CIFS), via Pietro Giuria 1, I-10125, Torino, Italy

<sup>6</sup> Dipartimento di Fisica "Ettore Pancini", Università di Napoli Federico II, via Cintia, 80126 Napoli, Italy

<sup>7</sup> INFN, Sezione di Napoli, via Cintia, 80126, Napoli, Italy

<sup>8</sup> Department of Physics and Astronomy, University of Southampton, Highfield, SO17 1BJ, UK.

January 14, 2019

## ABSTRACT

We present a catalog of 47 Wide-Angle Tailed radio galaxies (WATs), the WATCAT, selected combining observations from the National Radio Astronomy Observatory/Very Large Array Sky Survey (NVSS), the Faint Images of the Radio Sky at Twenty-Centimeters (FIRST) and the Sloan Digital Sky Survey (SDSS), and mainly built including a radio morphological classification. We included in the catalog only radio sources showing two-sided jets with two clear “warmspots” (i.e. jet knots as bright as 20% of the nucleus) lying on the opposite side of the radio core, and having classical extended emission resembling a plume beyond them. The catalog is limited to redshifts  $z \leq 0.15$ , and lists only sources with radio emission extended beyond 30 kpc from the host galaxy. We found that host galaxies of WATCAT sources are all luminous ( $-20.5 \gtrsim M_r \gtrsim -23.7$ ), red early-type galaxies with black hole masses in the range  $10^8 \lesssim M_{\text{BH}} \lesssim 10^9 M_{\odot}$ . The spectroscopic classification indicates that they are all low excitation galaxies (LEGs). Comparing WAT multifrequency properties with those of FR I and FR II radio galaxies at the same redshifts, we conclude that WATs show multifrequency properties remarkably similar to FR I radio galaxies, having radio power of typical FR IIs.

**Key words.** galaxies: active – galaxies: jets – radio continuum: galaxies

## 1. Introduction

We are now living in the golden age of wide and deep multifrequency surveys. This is extremely useful to investigate properties (see e.g., Kauffmann et al., 2003), nature, cosmological evolution (see e.g., Brandt and LSST AGN Science Collaboration, 2006; Best et al., 2014; Kaiser and Best, 2007) and environments (see e.g., Morganti et al., 1988; Croston et al., 2018; Hardcastle et al., 2018) of several classes of extragalactic sources, as radio-loud active galaxies, having a low sky density (i.e., number of sources per square degree). Thus, we recently embarked in a project aiming at creating homogeneous and complete catalogs of different classes of radio galaxies, uniformly selected on the basis of their radio morphological properties (Fanaroff & Riley, 1974). The first three catalogs, restricted to the local Universe (i.e., at  $z < 0.15$ ) and covering the major classes of radio galaxies (namely FR I, FR II and FR 0) have been recently published (Capetti et al., 2017a,b; Baldi et al., 2018), while the present work focuses again on a class of radio galaxies with extended radio emission beyond the host galaxy.

Both FR I and FR II radio galaxy catalogs were created on the basis of the radio morphological classification scheme of extragalactic radio sources with large-scale structures introduced by Fanaroff & Riley (1974). Thus radio galaxies can be broadly divided in two main classes, on the basis of the ratio of distance between the regions of highest radio surface brightness on opposite sides of the central nucleus (associated with the host galaxy) to the total extent of the source, up to the lowest brightness contour in the radio images at 1.4 GHz. Edge-darkened radio sources are classified as FR Is, while edge-brightened ones


as FR IIs. Then, in the FROCAT, we indeed presented a sample of 108 “compact” radio sources with  $z \leq 0.05$  and having a radio size  $\lesssim 5$  kpc and no extended radio emission. These also show the typical optical spectrum of low-excitation radio galaxies.

Here we present the fourth catalog of the series, focused on an “intermediate” class of radio galaxies between FR Is and FR IIs, having extended radio emission well beyond the optical profiles of their host galaxies: the Wide-Angle Tailed radio galaxies catalog.

Observed for the first time with the NRAO (National Radio Astronomy Observatory) interferometer at 2.7 GHz, during a survey of sources in the Abell clusters (Owen & Rudnick, 1976), WATs were named according to their radio morphology presenting bright hotspots (i.e., referred as “warmspots”) closer to their radio core with respect to FR IIs and with extended radio plumes beyond them (O’Donoghue et al., 1990).

Owen & Rudnick (1976), noting that this source class has higher radio luminosity than the radio sources with narrower tails (a.k.a. head-tailed radio galaxies), proposed that their curved radio structure could be due to the motion of the host galaxy through the intergalactic medium (IGM), slower than head-tailed sources. This scenario is supported by the fact that WATs are generally associated with more dominant cluster galaxies (Bird, 1994; Pinkney, 1995).

Sakelliou & Merrifield (2000) investigated the WAT radio structure using a sample selected from the those lying in Abell galaxy clusters. These authors suggested that, because the Universe evolves hierarchically with galaxy clusters forming through mergers of groups, WAT bent/curved tails are due to



**Fig. 1.** 1.4 GHz FIRST (black), 1.4 GHz NVSS (red) and 150 GHz TGSS (blue) radio contours of the WATCAT source SDSSJ130904.46+102935.3. Contours levels and increase factors are summarized in Table B.1. The field of view is  $3' \times 3'$ . The source name and redshift are reported in the upper right corner.

the ram pressure originated in merging processes. As pointed out by Hardcastle et al. (2005), similarly to FR IIs, WAT jets remain relativistic and well-collimated until they undergo rapid deceleration at a distance of few tens of kpc from the nucleus, creating the so called “warmspots”, the main feature of their radio morphology.

In FRI showing twin-jet (i.e., double sided), their well-collimated jet flow undergoes significant bulk deceleration on scales of a few kpc (Feretti et al., 1999; Mukherjee et al., 2013a,b; Laing and Bridle, 2014; Massaglia et al., 2016, 2018), slowing down from relativistic speeds to trans-sonic or sub-sonic regime with respect to the external medium (Laing and Bridle, 2002; Bicknell, 1994). This forms a jet that may in some cases make a smooth transition into a “plume” similar to those of observed in WATs (Leahy, 1993). Inner jets of FR Is are often also strong sources of X-ray emission (Hardcastle et al. 2001, 2003). WAT jets should be similar, and so we could expect to detect synchrotron X-ray emission arising from knots and “warmspots” (see also Massaro et al., 2011). It is worth noting that for all WATs being observed in the X-rays, as for FRI jets, the regions where their jets decelerate are coupled with high-energy particle acceleration that yield to X-ray synchrotron emission (e.g. Hardcastle et al. 2001; Worrall et al. 2001; Hardcastle et al. 2002). On the other hand, WAT plumes should not be similar to regions at the edges of FRI radio structures since the model outlined by Hardcastle & Sakellios (2004) requires the jet deceleration in the plumes to take place at a single discrete location.

Best et al. (2005), Baldi & Capetti (2010), and Best & Heckman (2012), to name recent examples, have analyzed properties of low redshift radio emitting AGNs using multifrequency information available thanks to NVSS, FIRST, SDSS. Here we just include radio morphological information to create the first catalog of WATs in the local Universe: hereinafter the WATCAT. The catalog we have built is the first to have this high level of completeness, and there are no other WATs catalogs with this population selected with so strict criteria.

WATs present a number of interesting problems and open questions, and tackling them can help us in achieving a better understanding of the dynamics of extragalactic radio sources and their interaction with the IGM.

This paper is organized as follows. In Sect. 2 we describe the selection criteria of sources included in the WATCAT, while radio and optical properties of the selected sources are then presented in Sect. 3. A comparison with FRI and FRII radio galaxies is discussed in Sect. 5 to dedicate Sect. 6 to our summary and conclusions.

A flat cosmology with  $H_0 = 67.8 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_M = 0.308$ , and  $\Omega_\Lambda = 0.692$  (Planck Collaboration et al., 2016) has been adopted throughout the paper. For our numerical results, we use c.g.s. units unless stated otherwise. Spectral indices  $\alpha$  are defined by the usual convention on the flux density,  $S_\nu \propto \nu^{-\alpha}$ . SDSS magnitudes are in the AB system and are corrected for the Galactic extinction. *WISE* magnitudes are instead in the Vega system and are not corrected for extinction since, as shown by, for example, D’Abrusco et al. (2014), such correction affects mainly the magnitude at  $3.4 \mu\text{m}$  of sources lying at low Galactic latitudes (and by less than  $\sim 3\%$ ). Not all WATs are detected in all three *WISE* bands centered at  $3.4$ ,  $4.6$  and  $12 \mu\text{m}$ , respectively. The detection rate is 78% at  $12 \mu\text{m}$  while it is 100% at  $3.4$  and  $4.6 \mu\text{m}$ .

## 2. Sample selection

We searched for WATs in the sample of 18,286 radio sources built by Best & Heckman, 2012 (hereafter BH12) by limiting our search to sources classified as active nuclei (AGN) on the basis of their radio emission. The BH12 catalog was created on the basis of observations available in the archive of the Sloan Digital Sky Survey (DR7/SDSS; Abazajian et al. 2009),<sup>1</sup> and combined with those of the National Radio Astronomy Observatory Very Large Array Sky Survey (NVSS; Condon et al. 1998) and the Faint Images of the Radio Sky at Twenty centimeters survey (FIRST; Becker et al. 1995) having the same footprint of the SDSS. For the radio data sets a radio flux density threshold of 5 mJy in the NVSS was adopted and the BH12 sample was also cross-matched with optical spectroscopic catalogs built by groups at the Max Planck Institute for Astrophysics and The Johns Hopkins University (Brinchmann et al., 2004; Tremonti et al., 2004).

As done in previous catalogs of FR Is and FR IIs, we selected 3,357 sources with redshift  $z \leq 0.15$ . Then, we visually inspected all FIRST images of each of these sources and selected only those whose radio emission is extended beyond 30 kpc from the center of the optical host galaxy, at the sensitivity of the FIRST. This cut led to a selected sample of 741 sources. We adopted a threshold of 30 kpc, mainly due to the limited resolution of FIRST survey, as estimate of the maximum size of the host galaxies. At redshift  $z = 0.15$  this distance corresponds to  $11''.4$ , thus ensuring that all 741 selected sources are well resolved with the  $5''$  resolution of the FIRST images. This allowed us to properly explore their morphology.

We chose a reference surface brightness level of 0.45 mJy/beam, correspondent to about three times the typical rms of FIRST images, at  $z = 0.15$ . As performed in previous radio galaxy catalogs we also corrected the radio surface brightness for the cosmological dimming and we applied a  $k$  correction by assuming a spectral index of 0.7, typical of lobes and plumes of radio galaxies.

Then, we selected WAT sources out of 741 objects on the basis of their radio morphology in FIRST images, considering strict criteria. Thus we included in the final sample only those radio sources showing two-sided jets with two clear “warmspots”

<sup>1</sup> <http://www.mpa-garching.mpg.de/SDSS/>.

(i.e. jet knots as bright as 20% of the nucleus) lying on the opposite side of the radio core and having classical extended emission resembling a plume beyond. Four authors performed this analysis independently and only the sources for which there was agreement by at least three of us were included in the final sample.

The final catalog of WATs (hereinafter the WATCAT) lists 47 sources, the first of this kind, homogenous at low redshifts.

In Fig. 1 we present the radio image with surface brightness contours from FIRST, NVSS and TIFR Giant Metrewave Radio Telescope (GMRT) Sky Survey (TGSS) radio maps of a typical WATCAT source. Images of all sources belonging to the WATCAT are available in the Appendix, while their main properties, as SDSS name, redshift, radio luminosity at 1.4 GHz and optical parameters are reported in Table 2. For each WAT, we provide both radio and [OIII] luminosities.

Then given the measurements of the velocity dispersions, available in the SDSS database, we also computed their black hole masses  $M_{BH}$  adopting the relation  $\sigma_* - M_{BH}$  of Tremaine et al. (2002). They range between  $7.9 \lesssim \log M_{BH} \lesssim 9.1 M_\odot$ , and their distribution peaks at  $\sim 10^{8.6} M_\odot$ . The uncertainty on the  $M_{BH}$ , of the order of a factor of 2, is mainly dominated by the dispersion of the correlation rather than by the uncertainty on  $\sigma_*$  measurements.

According to Montero-Dorta & Prada (2009) and the redshift completeness of the SDSS, the WATCAT, as the FRICAT and FRIICAT, is statistically complete at level of  $\sim 90\%$  in the optical band whereas this extremely low level of incompleteness is only due to a random loss of  $\sim 10\%$  of the potential spectroscopic targets (see, e.g., Zehavi et al., 2002). Therefore there could be the chance of missing radio sources featuring extended radio emission below the  $3\sigma$  threshold in FIRST images thus not able to resolve some diffuse emission.


On the other hand, to estimate the completeness of our catalog at radio frequencies, we re-scaled the luminosities of all sources in the  $z$  range between 0.05 and 0.09 computing their flux densities at 1.4 GHz and randomly assigning redshifts of sources in last  $z$  bin (see Fig. 2). Re-scaled flux densities all lie above the NVSS threshold. Then we also verified that the number of sources in the last  $z$  bin is consistent with the expected number extrapolated from the  $N \propto z^3$  scenario normalized to the first 4  $z$  bins. We found a good agreement as shown in Fig. 2. Consequently, we can assert that our sample is statistically complete in the radio band within the  $z$  range considered in our selection.

Finally we note that the BH12 catalog includes sources having the NVSS flux density at 1.4 GHz larger than 5 mJy while our selection is mainly based on the surface brightness distribution of FIRST images.

### 3. Multifrequency properties of WATs and their host galaxies

#### 3.1. WAT hosts: IR & optical properties

All optical spectra, obtained from the SDSS database, were inspected. We discovered that all sources minus one listed in the WATCAT are classified as low excitation galaxies (LEGs), in agreement with the classification reported by Best & Heckman (2012) based on the ratios of the optical emission lines in their SDSS spectra. The only exception is SDSS J155343.59+234825.4, being a high excitation galaxy (HEG). Given the high level of completeness of the WATCAT this is



**Fig. 2.** Distribution of the WATCAT sources redshifts. The blue line refers to the  $N \propto z^3$  relation normalized to the number of sources in the  $z$  bins between 0.05 and 0.09. The number of WATs in the  $z$  range between 0.13 and 0.15 appears to be consistent with this trend, indicating that the catalog is quite complete in the explored redshift range (see Section 2 for additional details).

the first time that is proven on a statistical basis for this source population that the majority of WATs is LEG.

The logarithm of [O III] line luminosities spans a range between 38.9 and 40.4 erg/s, with a mean value of 39.72 erg/s in log scale. However, it is worth noting that for four sources (SDSS J144904.27+025802.7, SDSS J141927.23+233810.2, SDSS J222455.24-002302.3, SDSS J155343.59+234825.4) we were not able to retrieve information regarding the [OIII] line luminosity, due to lack of this measurements in the SDSS database.


Then, thanks to the multifrequency observations available, as done in previous radio galaxy catalogs, we used various diagnostics to get a complete overview on the morphological and spectroscopic classification of the WAT host galaxies.

The distribution of absolute magnitude of the WATCAT host galaxies spans the range  $-20.5 \gtrsim M_r \gtrsim -23.7$  with a distribution peaking at  $M_r \sim -23$  (see Fig. 3).

The distribution of the concentration index  $C_r$ , defined as the ratio between the radii including 90% and 50% of the light in the  $r$  band, respectively, was obtained for each source directly from the SDSS database. The  $C_r$  distribution is consistent with WATs being hosted in early-type galaxies (ETGs). In general ETGs show higher values of  $C_r$  than late-type galaxies (LTGs). The thresholds adopted to distinguish ETGs and LTGs are the same used for the FRICAT and FRIICAT catalogs (see Strateva et al., 2001; Nakamura et al., 2003; Shen et al., 2003; Kauffmann et al., 2003; Bell et al., 2003; Bernardi et al., 2010, for more details).

We compared  $C_r$  values with those of the Dn(4000) spectroscopic index, defined as the ratio between the flux density measured on the “red” side of the Ca II break (4000–4100 Å) and that on the “blue” side (3850–3950 Å) (Balogh et al., 1999). Low redshift ( $z < 0.1$ ) red galaxies show  $Dn(4000) = 1.98 \pm 0.05$ , which is a value that decreases to  $= 1.95 \pm 0.05$  for  $0.1 < z < 0.15$  galaxies (Capetti & Raiteri, 2015). The presence of young stars or of nonstellar emission reduces the Dn(4000) index (Hamilton, 1985).

In Fig. 4 we show the concentration index  $C_r$  versus the Dn(4000) index (left panel) and versus  $M_{BH}$  (right panel) for



**Fig. 3.** Comparison between the normalized distributions of the host absolute magnitude in the  $r$  band ( $M_r$ ) for different classes of radio galaxies. The red histogram shows sources in the FRICAT, while the black one those listed in the FRIICAT, and the green one the selected WATCAT sources.


the WATCAT, FRICAT and FRIICAT sources. The vast majority of the WAT host galaxies lies in the region of high  $C_r$  and  $Dn(4000)$  values, indicating that they are all typical red ETGs. We performed a bidimensional Kolmogorov-Smirnov (KS) test Peacock (1983); Fasano and Franceschini (1987) and we found that comparing WATs with FR Is and FR IIs we obtain (1-p-value)  $\sim 10^{-16}$ , that implies that the two samples (WATs vs FR Is and WATs vs FR IIs) are not significantly different. There are only two exceptions, namely: SDSS J095716.41+190651.2 and SDSS J135315.36+550648.2 having both low values of the  $C_r$ . The distribution of black-hole masses ( see Fig. 5) shows that WATs, FR Is and FR II peak at the same value, but the FR IIs extend to lower values of the masses.

We also considered their  $u-r$  color while studying the properties of the WATCAT host galaxies. This allowed us to avoid uncertainties in the estimate of the  $Dn(4000)$  due to SDSS spectroscopic aperture, being of the order of  $3''$  in diameter. In Fig. 6 we show the  $u-r$  color versus the absolute  $r$ -band magnitude  $M_r$  of their hosts. WATs are all located above the line separating red and blue ETGs (consistently with their  $u-r$  distribution) thus following the same FR I trend (Schawinski et al., 2009). The fraction of “blue” ETGs decreases with increasing luminosity and these ETGs disappear for  $M_r \lesssim -22.5$ . The lack of blue ETGs among the WATCAT host galaxies is relevant.

The  $u-r$  distributions of WATs and FR IIs appear to be marginally different at a confidence level of 93% (p-value of 0.072). Same test performed for  $Dn(4000)$  index gave us a p-value of 0.58 for WATs and FR Is and 0.010 for WATs and FR IIs.

We note that, for the sake of clarity, in all our plots the uncertainties on optical parameters are not shown; however the average uncertainties are: 0.08, 0.03 and 0.004 on  $C_r$ ,  $Dn(4000)$  and  $m_r$ , respectively and  $\sim 9 \text{ km s}^{-1}$  on  $\sigma_*$ .

As last step, we cross-matched our WATCAT with the latest release of the Wide-field Infrared Survey Explorer (WISE) source catalog (i.e., AllWISE) retrieving mid-IR magnitudes. The associations between the WATCAT and the WISE catalog were computed adopting a  $3'3$  angular separation, which corresponds to the combination of the typical positional uncertainty




**Fig. 4.** (upper panel)  $Dn(4000)$  spectroscopic index vs. concentration index  $C_r$  for WATCAT sources (green diamonds), FRICAT sources (black dots), and FRIICAT sources (red squares). It appears that WATs are hosted in red early type galaxies (ETGs) like FR Is, while high excitation (HEG) FR IIs show lower values of  $Dn(4000)$ . (lower panel) Concentration index  $C_r$  vs black hole mass (in solar units). WATs show a similar behaviour to FR Is, while FR IIs extend to lower black hole masses with respect to the WATCAT sample, probably due to relatively large errors particularly in the measurement of  $\sigma_*$  a possible uncertain identification of their spectroscopic class, or a substantial contribution from a bright non-thermal nucleus.


of the WISE all sky survey (Wright et al., 2010) and that of the FIRST (D’Abrusco et al., 2014). In Fig. 7 we show the comparison between the mid-IR colors of WATCAT, FRICAT and FRIICAT sources.

WATCAT sources appear to have WISE colors mostly dominated by their host galaxies (they fall in the same region as elliptical galaxies; Wright et al. 2010) and not contaminated by the non-thermal emission of their jets, as for example occurs in blazars (D’Abrusco et al., 2012; Massaro et al., 2011; Massaro & D’Abrusco, 2016). Only one source (SDSS J155343.59+234825.4), classified as HEG, has  $w_2 - w_3 > 0.3$  and it is located at the onset of the sequence defined by the more luminous objects such as blazars and flat spectrum radio quasars (Massaro et al., 2012). Thus, the WISE infrared colors further support the idea that WATs are hosted in “passive” ETGs.





**Fig. 5.** Comparison between black holes masses distributions for WATCAT, FRICAT and FRIICAT hosts.




**Fig. 6.**  $u-r$  color vs. absolute  $r$  band magnitude,  $M_r$ , for the WATCAT FRICAT and FRIICAT hosts (the green diamond represent WATs, the black dots represent the FR Is sample and the red squares the FR IIs). WATs are associated with red ETGs. The dashed blue line represents the relation from Schawinski et al. (2009), that separates red and blue ETGs.


### 3.2. Radio properties

In Fig. 8, we show the distribution of the NVSS radio luminosity at 1.4 GHz (i.e.,  $\nu L_\nu$ ) for all the sources belonging to the WATCAT in comparison with FR I and FR II listed in Capetti et al. (2017a,b). The distribution of radio luminosity at 1.4 GHz of the WATCAT covers the range  $L_{1.4} = \nu_r L_r = \sim 10^{40.1} - 10^{41.6}$  erg s<sup>-1</sup>. The Fanaroff & Riley (1974) separation between FR Is and FR IIs translates, with our adopted cosmology and by assuming a spectral index of 0.7 between 178 MHz and 1.4 GHz, into  $L_{1.4} \sim 10^{41.6}$  erg s<sup>-1</sup>.

The distribution of the NVSS radio luminosity at 1.4 GHz clearly shows that WATs are on average more powerful than FR Is and have radio luminosity similar to classical FR II radio




**Fig. 7.** WISE mid-IR colors of the WATCAT hosts (green diamonds) compared to those of FRICAT (black dots) and FRIICAT (red squares). WATCAT sources appear to have mid-IR colors mostly dominated by their host galaxies because they lie in the same color space region as elliptical galaxies, and do not show contamination from the emission of their jets.



**Fig. 8.** Normalized distribution of the NVSS luminosities at 1.4 GHz. The black histogram represents the 219 FRICAT sources, the green histogram the 47 WATCAT sources, and the red histogram the 122 FRIICAT sources. WATs are on average more powerful than FR Is and have radio luminosity similar to classical FR II radio galaxies.

galaxies. According to a simple KS test we found WATs and FR IIs are indistinguishable while for the comparison between WATs and FR Is these two population differs within a confidence level of  $\sim 10^{-15}$ .

We measured the angular separation between the peaks of surface brightness of two “warmspots” for each WAT on the FIRST radio maps at 1.4 GHz. Then we converted the angular separation into physical distance in kpc, so defining their linear size:  $d_w$ . In Fig. 9 we show the relation between the  $d_w$  and the WAT radio power. The low number of WATs, coupled with functional biases in the relations used to compute  $d_w$  and the radio power, prevents us from evaluating a correlation between these two parameters, however there is a hint that high radio power



**Fig. 9.** NVSS radio luminosity at 1.4 GHz vs linear size, that is physical separation between the “warmspots”. We note that at large physical separations “warmspots” of low radio luminosity are not produced, i.e., the region in the upper left corner appears empty.

WATs do not produce “warmspots” closer to the radio core with respect to low power ones is present.

#### 4. Discussion

Here we discuss the multifrequency properties of the WATCAT in comparison with both the FR Is and FR IIs listed in the FRICAT and FRIICAT.

Host galaxies of WATCAT sources are remarkably uniform being all luminous red ETGs and spectroscopically classified as LEGs, as occurs for the FR Is. On the other hand, their radio luminosity at 1.4 GHz is generally higher than that of FR Is, regardless of the value of  $L_{\text{[O III]}}$ .

A KS test to compare the  $r$  band absolute magnitude distributions of WATs, FR Is and FR IIs was performed. We found that WATs and FR Is distributions of  $M_r$  differ while this difference is only marginal between WATs and FR IIs (i.e.,  $p$ -value of 0.013). This can be interpreted according to the scenario where WATs tend to be the brightest cluster galaxy when hosted in massive galaxy clusters, while this is not always true for the FR Is (Quintana and Lawrie, 1982).

In 1974, Fanaroff & Riley also observed that all sources in their original sample with luminosity measured at 178 MHz lower than  $2 \times 10^{25}$  W/Hz/sr (assuming a Hubble constant of  $50 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ) were classified as FR Is, while more powerful radio objects were all presenting the FR II morphology (Fanaroff & Riley, 1974). This dichotomy between FR classes is fairly sharp at low radio frequencies (see also ), but Ledlow & Owen (1996) showed that their separation is clearer when comparing radio powers with optical magnitudes of their host galaxies.

As previously carried out for the FRICAT and FRIICAT samples, we built a re-scaled diagram plotting the radio power vs the optical magnitudes for our WATs. The bulk of WATCAT sources lie below the boundary between FRI and FR II radio galaxies as shown in Fig. 10), mostly lying in the region populated by FRI sources. We remark that the dividing line between the two main classes of radio galaxies highlighted by Ledlow and Owen (1976) was appropriately shifted including a correction of 0.12 mag to the magnitude of the host galaxies and a correction of 0.22 mag to convert the Cousin magnitude system into the SDSS

one (Fukugita et al., 1996). The location of WATs in this radio power vs optical magnitude confirms that the more powerful WATs tend to be associated with more massive galaxies.

As previously shown, classical FR Is belonging to the 3C sample show a positive trend between the [OIII] emission line and radio luminosity with a slope consistent with unity (e.g., Buttiglione et al. 2010). This indicates that a constant fraction of the AGN power, as measured from the  $L_{\text{[O III]}}$ , is then converted into radio power. Buttiglione et al. show that the same result, although with a different normalization, is found when considering the FR II radio galaxies again selected out of the 3C.

No neat trend or correlation between  $L_{1.4}$  and  $L_{\text{[O III]}}$  can be seen for WATs, as occurs for FR Is, (see Figure 11) where, at a given emission line luminosity, radio power spans over two orders of magnitude. Similarly to 3C radio galaxies, it appears that no source has  $aL_{1.4}/L_{\text{[O III]}}$  ratio exceeding  $\sim 100$ , producing the scarcely populated region in the bottom right portion of this plot. On the other hand several WATs show values of the  $L_{1.4}/L_{\text{[O III]}}$  ratio below the 0.5 threshold. The broad distribution of the  $L_{1.4}/L_{\text{[O III]}}$  ratio points that a broad range AGN bolometric power is then converted into radio emission. Furthermore, radio luminosity increases, not surprisingly, when the size of the radio source increases.

Finally, we remark that, as occurred for FR Is both spectroscopic and host properties WAT host galaxies rule out the possibility of having the [OIII] emission line contaminated by star formation thus, we would have been able to observe a trend, if present between radio power and emission line luminosity of the [OIII].

#### 5. Summary and conclusions


We built a low redshift (i.e.  $z \leq 0.15$ ) catalog of 47 WATs, the WATCAT. This sample was selected out of the Best & Heckman (2012) catalog, built combining observations available in the SDSS, NVSS, and FIRST surveys, but including a radio morphological classification.

The WAT classification is purely morphological and based on the visual inspection of FIRST radio images at 1.4 GHz. We included sources in which the radio emission extends beyond 30 kpc from the position of their optical host galaxy. Our final catalog is also restricted to redshift  $z < 0.15$ .


WATs show radio jets, as in FR IIs, that are thought to remain relativistic and well-collimated. WATs are radio sources showing two-sided jets with two clear “warmspots” (i.e. jet knots as bright as 20% of the nucleus) lying on the opposite side of the radio core, and having classical extended emission resembling a plume beyond them.

Our main results are listed below:

- Using NVSS radio observations at 1.4 GHz, we found out that WATs tend to be more radio luminous than FR Is, and more similar to FR IIs;
- WATCAT host galaxies are remarkably homogeneous, as they do not show significant differences in their optical and infrared colors with respect to the general population of massive ETGs and the presence of an active nucleus (and its level of activity) does not appear to affect their host galaxies;
- SDSS optical spectra of WATCAT sources proved that these sources are all classified as low excitation radio galaxies (LEGs). FR Is are all classified as LEGs, while FR IIs can be LEGs or high excitation radio galaxies (HEGs). This again marks a similar behavior between WATs and FR Is;



**Fig. 10.** (left panel): Host absolute magnitude ( $M_r$ ) vs. radio luminosity (NVSS) at 1.4 GHz, for WATCAT, FRICAT and FRIICAT sources (green diamonds, black dots and red squares, respectively). The dashed blue line shows the separation between FR I and FR II as reported by Ledlow & Owen (1996) to which we applied a correction of 0.34 mag to account for the different magnitude definition and the color transformation between the SDSS and Cousins systems. The bulk of the WATCAT sources lies below the boundary between FR I and FR II radio galaxies. Right panel: radio luminosity (NVSS) at 1.4 GHz vs. black hole mass for WATCAT, FRICAT and FRIICAT (green diamonds, black dots and red squares, respectively). WATs and FR Is show similar black holes masses, but WATs are more radio luminous.



**Fig. 11.** Comparison between [O III] line luminosity and radio luminosity (NVSS) of the WATCAT, FRICAT and FRIICAT samples (green diamonds, black dots and red squares, respectively). The turquoise line shows the linear correlation between these two quantities derived from the FR Is of the 3C sample (Buttiglione et al., 2010). No correlation between  $vL_v$  and  $L_{[O III]}$  can be seen for the WATCAT (as for FRICAT) where, at a given line luminosity, the radio luminosities span over 1-1.5 orders of magnitude.

- $M_r$  vs  $vL_v$  measured at 1.4 GHz showed that WATs follow the same behavior of FR II HEGs;
- using infrared data collected by *WISE* WAT host galaxies appear to be similar to normal elliptical galaxies, with IR colors similar to those of FR Is, lacking significant contribution from active star formation;
- values of the absolute magnitude in the r band ( $M_r$ ) are also in agreement with the fact that WATs, as reported in litera-

ture, could be the brightest cluster galaxy when they resides therein;

- comparing the distribution of the concentration index (i.e.,  $C_r$ ) and the dispersion index  $Dn(4000)$ , we discovered that WATs tend to not be extremely “blue” as occurs for FR Is, lacking signatures of active star formation, that are sometimes found in FR IIs;
- from the comparison between the radio luminosity at 1.4 GHz, obtained from NVSS, and the luminosity of the [OIII] emission line present in WAT optical spectra, we found that our selected WATs do not follow any trend between  $L_{1.4}$  and  $L_{[O III]}$  having [OIII] emission line luminosity even lower than expected for a given value of the radio power at 1.4 GHz.

Finally, we conclude that WATs show multifrequency properties remarkably similar to FRI radio galaxies, being more powerful at radio frequencies and similar to typical FR IIs.

## References

- Abazajian, K. N., Adelman-McCarthy, J. K., Agüeros, M. A., et al. 2009, *ApJS*, 182, 543
- Baldi, R. D. & Capetti, A. 2010, *A&A*, 519, A48+
- Baldi, R. D., Capetti, A., Massaro, F. 2018, *A&A*, 609, A1.
- Balogh, M. L., Morris, S. L., Yee, H. K. C., Carlberg, R. G., & Ellingson, E. 1999, *ApJ*, 527, 54
- Becker, R. H., White, R. L., & Helfand, D. J. 1995, *ApJ*, 450, 559
- Bell, E. F., McIntosh, D. H., Katz, N., & Weinberg, M. D. 2003, *ApJS*, 149, 289
- Bernardi, M., Shankar, F., Hyde, J. B., et al. 2010, *MNRAS*, 404, 2087
- Best, P. N. & Heckman, T. M. 2012, *MNRAS*, 421, 1569
- Best, P. N., Kauffmann, G., Heckman, T. M., et al. 2005, *MNRAS*, 362, 25
- Best, P. N., Ker, L. M., Simpson, C., Rigby, E. E., Sabater, J. 2014, *MNRAS*, 445, 955.
- Bicknell, G. V. 1994, *Australian Journal of Physics*, 47, 669.
- Brandt, N., LSST AGN Science Collaboration 2006. AGN Science with the LSST. American Astronomical Society Meeting Abstracts 209, 86.14
- Bird, C. M. 1994. Substructure in Clusters and Central Galaxy Peculiar Velocities. *The Astronomical Journal* 107, 1637.
- Brinchmann, J., Charlot, S., White, S. D. M., et al. 2004, *MNRAS*, 351, 1151
- Burns, J. O. 1981, *MNRAS*, 195, 523.

- Buttiglione, S., Capetti, A., Celotti, A., et al. 2010, *A&A*, 509, A6
- Capetti, A. & Raiteri, C. M. 2015, *A&A*, 580, A73
- Capetti, A., Massaro, F., Baldi, R. D. 2017, *A&A* 598, A49.
- Capetti, A., Massaro, F., Baldi, R. D. 2017, *A&A*, 601, A81.
- Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, *AJ*, 115, 1693
- Croston, J. H., and 15 colleagues 2018, *ArXiv e-prints arXiv:1811.07949*.
- D'Abrusco, R., Massaro, F., Ajello, M., et al. 2012, *ApJ*, 748, 68
- D'Abrusco, R., Massaro, F., Paggi, A., et al. 2014, *ApJS*, 215, 14
- Edge, D. O., Shakeshaft, J. R., McAdam, W. B., Baldwin, J. E., Archer, S. 1959. A survey of radio sources at a frequency of 159 Mc/s.. *Memoirs of the Royal Astronomical Society* 68, 37.
- Eilek, J. A., Henriksen, R. N. 1984, *AJ*, 277, 820.
- Fanaroff, B. L. & Riley, J. M. 1974, *MNRAS*, 167, 31P
- Fasano, G., Franceschini, A. 1987. A multidimensional version of the Kolmogorov-Smirnov test. *Monthly Notices of the Royal Astronomical Society* 225, 155.
- Feretti, L., Perley, R., Giovannini, G., Andernach, H. 1999., *Memorie della Societa Astronomica Italiana* 70, 129.
- Fukugita, M., Ichikawa, T., Gunn, J. E., et al. 1996, *AJ*, 111, 1748
- Hamilton, D. 1985, *AJ*, 297, 371.
- Hardcastle, M. J., Alexander, P., Pooley, G. G., Riley, J. M. 1998, *MNRAS*, 296, 445
- Hardcastle, M. J., Birkinshaw, M., Worrall, D. M. 2001, *MNRAS*, 323, L17.
- Hardcastle, M. J., Worrall, D. M., Birkinshaw, M., Laing, R. A., Bridle, A. H. 2002, *MNRAS*, 334, 182.
- Hardcastle, M. J., Sakelliou, I. 2004, *MNRAS*, 349, 560.
- Hardcastle, M. J., Sakelliou, I., Worrall, D. M. 2005, *MNRAS*, 359, 1007.
- Hardcastle, M. J., and 22 colleagues 2018. Radio-loud AGN in the first LoTSS data release: The lifetimes and environmental impact of jet-driven sources. *arXiv e-prints arXiv:1811.07943*.
- Kaiser, C. R., Best, P. N. 2007, *MNRAS*, 381, 1548.
- Kauffmann, G., Heckman, T. M., White, S. D. M., et al. 2003, *MNRAS*, 341, 33
- Laing, R. A., Bridle, A. H. 2002, *MNRAS*, 336, 1161.
- Laing, R. A., Bridle, A. H. 2014, *MNRAS*, 437, 3405.
- Leahy, J. P. 1993. DRAGNs. *Jets in Extragalactic Radio Sources* 1.
- Ledlow, M. J. & Owen, F. N. 1996, *AJ*, 112, 9
- Massaglia, S., Bodo, G., Rossi, P., Capetti, S., Mignone, A. 2016, *A&A*, 596, A12.
- Massaglia, S., Bodo, G., Rossi, P., Capetti, S., Mignone, A. 2018, *ArXiv e-prints arXiv:1812.00657*.
- Massaro, F., D'Abrusco, R., Ajello, M., Grindlay, J. E., & Smith, H. A. 2011, *ApJ*, 740, L48
- Massaro, F., D'Abrusco, R. 2016. The Infrared-Gamma-Ray Connection: A WISE View of the Extragalactic Gamma-Ray Sky. *The Astrophysical Journal* 827, 67.
- Massaro, F., Harris, D. E., Cheung, C. C. 2011. Large-scale Extragalactic Jets in the Chandra Era. I. Data Reduction and Analysis. *The Astrophysical Journal Supplement Series* 197, 24.
- Massaro, F., D'Abrusco, R., Tosti, G., et al. 2012, *ApJ*, 750, 138
- Montero-Dorta, A. D. & Prada, F. 2009, *MNRAS*, 399, 1106
- Morganti, R., Fanti, R., Gioia, I. M., Harris, D. E., Parma, P., de Ruiter, H. 1988. Low luminosity radio galaxies : effects of gaseous environment.. *Astronomy and Astrophysics* 189, 11.
- Mukherjee, D., Bhattacharya, D., Mignone, A. 2013. MHD instabilities in accretion mounds - I. 2D axisymmetric simulations. *Monthly Notices of the Royal Astronomical Society* 430, 1976.
- Mukherjee, D., Bhattacharya, D., Mignone, A. 2013. MHD instabilities in accretion mounds - II. 3D simulations. *Monthly Notices of the Royal Astronomical Society* 435, 718
- Nakamura, O., Fukugita, M., Yasuda, N., et al. 2003, *AJ*, 125, 1682
- O'Donoghue, A. A., Owen, F. N., Eilek, J. A. 1990, *ApJS*, 72, 75.
- Owen, F. N. & Rudnick, L. 1976, *ApJ*, 205, L1
- Peacock, J. A. 1983. Two-dimensional goodness-of-fit testing in astronomy.. *Monthly Notices of the Royal Astronomical Society* 202, 615.
- Pinkney, J. 1995. The Dynamics of Galaxy Clusters Containing Wide - Tailed Radio Sources.. *Ph.D. Thesis*.
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, *A&A*, 594, A13
- Quintana, H., Lawrie, D. G. 1982. On the determination of velocity dispersions for cD clusters of galaxies.. *The Astronomical Journal* 87, 1.
- Rudnick, L. & Owen, F. N. 1977, *AJ*, 82, 1
- Sakelliou, I., Merrifield, M. R. 2000, *MNRAS*, 311, 649.
- Schawinski, K., Lintott, C., Thomas, D., et al. 2009, *MNRAS*, 396, 818
- Shen, S., Mo, H. J., White, S. D. M., et al. 2003, *MNRAS*, 343, 978
- Strateva, I., Ivezić, Ž., Knapp, G. R., et al. 2001, *AJ*, 122, 1861
- Tremaine, S., Gebhardt, K., Bender, R., et al. 2002, *ApJ*, 574, 740
- Tremonti, C. A., Heckman, T. M., Kauffmann, G., et al. 2004, *ApJ*, 613, 898
- Worrall, D. M., Birkinshaw, M., Hardcastle, M. J. 2001, *MNRAS*, 326, L7.
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868
- Zehavi, I., Blanton, M. R., Frieman, J. A., et al. 2002, *ApJ*, 571, 172
- Zirbel, E. L. & Baum, S. A. 1995, *ApJ*, 448, 521

*Acknowledgements.* This work is supported by the "Departments of Excellence 2018 - 2022" Grant awarded by the Italian Ministry of Education, University and Research (MIUR) (L. 232/2016). This research has made use of resources provided by the Compagnia di San Paolo for the grant awarded on the BLENV project (S1618L1.MASF.01) and by the Ministry of Education, Universities and Research for the grant MASF.FFABR.17.01. This investigation is supported by the National Aeronautics and Space Administration (NASA) grants GO4-15096X, AR6-17012X and GO6-17081X. F.M. acknowledges financial contribution from the agreement ASI-INAf n.2017-14-H.0 A.P. acknowledges financial support from the Consorzio Interuniversitario per la fisica Spaziale (CFIS) under the agreement related to the grant MASF.CONTR.FIN.18.02. Part of this work is based on the NVSS (NRAO VLA Sky Survey): The National Radio Astronomy Observatory is operated by Associated Universities, Inc., under contract with the National Science Foundation. This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration. This research made use of the NASA/IPAC Infrared Science Archive and Extragalactic Database (NED), which are operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. Funding for SDSS-III has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, and the U.S. Department of Energy Office of Science. The SDSS-III web site is <http://www.sdss3.org/>. SDSS-III is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS-III Collaboration, including the University of Arizona, the Brazilian Participation Group, Brookhaven National Laboratory, University of Cambridge, Carnegie Mellon University, University of Florida, the French Participation Group, the German Participation Group, Harvard University, the Instituto de Astrofísica de Canarias, the Michigan State/Notre Dame/JINA Participation Group, Johns Hopkins University, Lawrence Berkeley National Laboratory, Max Planck Institute for Astrophysics, Max Planck Institute for Extraterrestrial Physics, New Mexico State University, New York University, Ohio State University, Pennsylvania State University, University of Portsmouth, Princeton University, the Spanish Participation Group, University of Tokyo, University of Utah, Vanderbilt University, University of Virginia, University of Washington, and Yale University.



Source name	Low contour level FIRST	Contour increase factor	Low contour level NVSS	Contour increase factor	Low contour level TGSS	Contour increase factor
J004312.85-103956.1	0.001	x2	0.001	x4	0.01	x2
J080101.35+134952.1	0.001	+2	0.002	x2	0.04	+2
J080337.67+105042.4	0.001	+2	0.01	x2	0.02	+2
J081803.86+543708.4	0.001	x2	0.001	+2	0.01	x2
J082718.31+463510.9	0.001	+2	0.001	x4	0.01	x2
J085116.23+082723.1	0.001	+2	0.001	x2	0.01	x2
J091337.21+031720.5	0.001	x2	0.001	x4	0.002	x2
J092428.89+141409.3	0.002	x2	0.002	x2	0.02	x4
J092539.05+362705.5	0.001	+2	0.001	x2	0.02	x2
J092612.34+324721.2	6.0E-4	+2	0.002	+2	0.01	x2
J093349.82+451957.8	4.0E-4	x2	0.001	x2	0.008	+4
J095716.40+190651.2	0.001	+2	0.001	x2	0.01	x2
J101932.33+140301.8	0.001	x2	0.001	x2	0.01	x2
J103502.61+425548.3	0.001	x2	0.001	x2	0.008	x2
J103605.76+000606.8	0.001	x2	0.001	x4	0.02	x2
J103636.24+383508.1	0.001	+2	0.001	+2	0.01	+2
J103856.37+575247.5	0.001	+2	0.001	x2	0.02	x2
J104645.86+314426.8	0.001	+2	0.001	x2	0.01	+2
J104914.08+005945.2	0.001	+2	0.002	x2	0.01	+2
J114020.22+535029.1	0.001	+2	0.001	+2	0.006	x2
J114111.81+054404.9	0.001	x2	0.002	x2	0.01	x2
J115424.56+020653.0	0.001	+2	0.001	+2	0.01	+2
J115513.65-003133.9	0.001	x2	0.002	+2	0.01	+2
J120118.19+061859.3	0.001	x2	0.001	x2	0.01	x2
J120455.01+483256.9	0.001	x2	0.002	x2	0.01	x2
J121439.53+052803.9	0.001	x2	0.001	x4	0.01	x2
J130904.46+102935.3	0.001	x2	0.002	x2	0.01	x4
J133038.38+381609.7	0.001	x2	0.001	x4	0.01	x4
J135315.36+550648.3	0.001	+2	0.001	x2	0.01	x4
J141456.58+001223.0	0.001	x2	0.002	x2	0.01	x2
J141513.98-013703.7	0.001	x2	0.001	x2	0.01	x2
J141718.94+060812.3	0.001	x2	0.002	x2	0.08	x2
J141731.27+081230.1	0.001	x2	0.001	x4	0.04	x2
J141927.23+233810.2	4.0E-4	x2	0.001	x2	0.01	x2
J143304.34+033037.6	0.001	x2	0.001	x2	0.01	x2
J143409.03+013700.9	0.001	x2	0.001	x4	0.01	x2
J144700.45+460243.5	0.001	x2	0.001	x2	0.01	x2
J144904.27+025802.7	0.001	+1	0.001	x2	0.02	x2
J150229.04+524402.1	0.001	x2	0.001	x2	0.02	x2
J151108.78+180153.2	0.001	x2	0.002	x4	0.02	x2
J154346.14+341521.6	4.0E-4	x2	0.002	x2	0.01	x2
J154729.58+145656.9	0.001	x2	0.002	x2	0.02	x2
J155343.59+234825.4	0.001	x4	0.002	x2	0.01	x2
J161828.98+295859.5	0.001	x2	0.001	x2	0.01	x2
J164527.68+272005.8	0.001	x2	0.001	x2	0.01	x2
J212546.35+005551.9	0.001	x2	0.001	x4	0.01	x2
J222455.24-002302.3	0.001	x2	0.002	x2	0.01	x2

**Table 1.** Column description: (1) source name; (2) lower FIRST contour level (mJy); (3) contour increase factor (mJy); (4) lower NVSS contour level (mJy); (5) contour increase factor (mJy); (6) lower TGSS contour level (mJy); (7) contour increase factor (mJy).

Source name	$z$	$\log \nu L_\nu$	$M_r$	$\log L[\text{OIII}]$	Dn	$C_r$	$\sigma^*$
J004312.85-103956.0	0.13	40.93	-23.50	19.85	1.93	3.18	290.93
J080101.35+134952.2	0.11	41.21	-23.32	22.18	1.95	3.15	221.46
J080337.67+105042.4	0.14	40.98	-22.18	3.06	2.00	3.49	246.58
J081803.86+543708.4	0.12	41.43	-23.67	22.72	1.96	3.09	263.64
J082718.31+463510.8	0.12	40.74	-22.60	13.73	2.01	3.34	277.60
J085116.24+082723.1	0.06	40.36	-22.40	51.05	1.97	3.22	258.04
J091337.21+031720.5	0.14	40.94	-22.50	6.98	1.78	2.57	236.35
J092428.89+141409.3	0.14	40.44	-22.71	11.58	1.98	3.29	242.06
J092539.06+362705.6	0.11	41.49	-23.48	26.98	1.97	3.28	276.34
J092612.34+324721.2	0.14	41.38	-21.97	14.35	1.82	2.88	179.16
J093349.82+451957.8	0.13	41.33	-23.05	20.75	1.90	2.94	218.17
J095716.41+190651.2	0.09	40.88	-22.17	35.13	1.91	1.42	210.85
J101932.33+140301.8	0.15	41.33	-23.21	15.80	1.97	2.92	264.37
J103502.62+425548.3	0.14	40.86	-22.88	17.15	2.22	3.19	251.18
J103605.76+000606.8	0.10	41.25	-23.32	39.53	1.88	3.03	311.06
J103636.24+383508.1	0.14	41.14	-22.84	18.27	2.03	3.20	280.19
J103856.37+575247.5	0.10	40.95	-21.43	20.40	1.92	2.93	210.19
J104645.86+314426.8	0.11	40.81	-23.02	19.30	2.01	3.22	278.41
J104914.08+005945.2	0.11	40.63	-22.78	12.53	1.91	3.19	266.85
J114020.23+535029.1	0.15	41.04	-23.17	11.86	2.09	3.05	332.40
J114111.81+054405.0	0.10	41.24	-23.47	1.06	1.93	2.94	272.32
J115424.56+020653.0	0.13	40.64	-22.90	5.80	1.97	3.52	276.32
J115513.65-003133.9	0.13	41.02	-22.76	17.40	1.92	2.85	221.82
J120118.19+061859.3	0.14	40.54	-22.81	8.39	1.99	3.32	319.14
J120455.02+483256.9	0.07	40.62	-21.81	26.41	1.90	3.23	233.62
J121439.53+052803.9	0.08	40.13	-23.44	5.01	1.95	3.28	230.03
J130904.46+102935.3	0.09	41.17	-22.87	4.74	2.02	3.44	348.60
J133038.38+381609.7	0.11	40.86	-23.24	24.97	1.94	3.17	260.61
J135315.36+550648.2	0.14	40.66	-20.49	6.37	1.96	2.01	255.58
J141456.58+001223.0	0.13	40.82	-22.87	15.15	2.01	3.28	331.01
J141513.98-013703.7	0.15	41.24	-23.33	37.24	2.00	3.05	305.79
J141718.94+060812.3	0.11	40.91	-22.79	2.55	2.00	3.31	278.47
J141731.27+081230.1	0.06	40.58	-22.52	49.76	2.04	3.33	237.39
J141927.23+233810.2	0.14	40.12	-22.16	-1.67	1.88	2.94	185.40
J143304.34+033037.6	0.15	40.88	-23.44	17.21	1.98	2.65	230.86
J143409.03+013700.9	0.14	41.61	-23.43	18.35	2.01	3.35	331.87
J144700.45+460243.5	0.13	41.04	-22.34	18.54	1.92	3.36	288.44
J144904.27+025802.7	0.12	40.36	-21.81	-3.85	1.87	3.32	194.29
J150229.04+524402.0	0.13	41.10	-23.50	24.43	1.98	3.28	292.56
J151108.77+180153.3	0.12	41.28	-23.08	21.10	2.04	3.11	274.49
J154346.14+341521.6	0.12	40.20	-22.30	3.30	1.93	3.29	232.29
J154729.59+145657.0	0.09	40.69	-22.42	37.78	1.85	3.02	203.33
J155343.59+234825.4	0.12	41.28	-23.43	307.83	1.64	3.18	323.50
J161828.98+295859.6	0.13	40.96	-23.21	5.99	2.04	3.15	315.10
J164527.68+272005.8	0.10	40.65	-23.16	14.75	1.96	3.05	270.40
J212546.35+005551.8	0.14	41.62	-23.02	25.49	1.96	2.98	223.22
J222455.24-002302.3	0.14	40.74	-22.63		2.7		274.29


**Table 2.** Column description: (1) source name; (2) redshift; (3) logarithm of the radio luminosity [ $\text{erg s}^{-1}$ ]; (4) SDSS DR7 r band AB magnitude; (5) logarithm of the [O III] line luminosity [ $\text{erg s}^{-1}$ ] (6) Dn(4000) spectroscopic index; (7) concentration index  $C_r$ ; (8) stellar velocity dispersion [ $\text{km s}^{-1}$ ].

Best ID	SDSS name
84	J103605.76+000606.8
101	J104914.08+005945.2
166	J115513.65-003133.9
383	J141456.58+001223.0
1916	J115424.56+020653.0
2105	J143409.03+013700.9
2132	J144904.27+025802.7
2241	J082718.31+463510.8
2412	J091337.21+031720.5
2630	J143304.34+033037.6
3222	J004312.85-103956.0
4363	J114111.81+054405.0
4409	J121439.53+052803.9
5113	J141513.98-013703.7
5428	J103856.37+575247.5
5754	J212546.35+005551.8
6044	J114020.23+535029.1
6456	J150229.04+524402.0
6781	J093349.82+451957.8
7294	J092539.06+362705.6
7770	J135315.36+550648.2
8653	J154346.14+341521.6
8775	J161828.98+295859.6
8847	J103502.62+425548.3
8860	J103636.24+383508.1
9093	J120455.02+483256.9
9641	J092612.34+324721.2
9939	J120118.19+061859.3
10259	J144700.45+460243.5
10402	J164527.68+272005.8
11011	J101932.33+140301.8
11140	J085116.24+082723.1
11351	J081803.86+543708.4
11515	J130904.46+102935.3
11671	J141731.27+081230.1
11817	J141718.94+060812.3
13055	J133038.38+381609.7
13211	J104645.86+314426.8
13950	J141927.23+233810.2
14994	J080101.35+134952.2
15652	J095716.41+190651.2
15725	J080337.67+105042.4
15953	J092428.89+141409.3
16471	J154729.59+145657.0
858	J222455.24-002302.3
12051	J155343.59+234825.4
17907	J151108.77+180153.3

**Table A.1.** Column description: (1) Best ID, from BH12; (2) SDSS DR9 source name.

## Appendix A: FIRST radio contours of the 47 WATCAT sources.

In this Appendix we show FIRST radio contours for the 47 sources in the WATCAT.



**Fig. A.1.** Images of the WATCAT sources, ordered by right ascension. Contours, corrected for cosmological effects to a redshift of  $z = 0.15$ , are drawn starting from 0.45 mJy/beam and increase with a geometric progression with a common ratio of  $\sqrt{2}$ . The field of view is  $3' \times 3'$ . The black circle is centered on the host galaxy and has a radius of 30 kpc. The source ID and redshift are reported in the upper corners.






Fig. A.1. (continued)




Fig. A.1. (continued)




Fig. A.1. (continued)





Fig. A.1. (continued)



**Appendix B: FIRST, NVSS and TGSS radio contours of the 47 *WATCAT* sources.**

In this appendix we show FIRST, NVSS and TGSS radio contours for the 47 sources in the *WATCAT*.



**Fig. B.1.** Images of the WATCAT sources, ordered by right ascension. See Table for contours level. The field of view is  $3' \times 3'$ ; the black tick at the bottom is 150 kpc long. The source name and redshift are reported in the upper right corner.




Fig. B.1. (continued)




Fig. B.1. (continued)






Fig. B.1. (continued)